arXiv:1702.08023v1 [cs.FL] 26 Feb 2017

A SURVEY ON DIFFERENCE HIERARCHIES OF REGULAR LANGUAGES

OLIVIER CARTON, DOMINIQUE PERRIN, AND JEAN-ÉRIC PIN

IRIF, CNRS and Université Paris-Diderot,, Case 7014, 75205 Paris Cedex 13, France.
e-mail address: olivier.carton@irif.fr

Laboratoire d'informatique Gaspard-Monge, Université de Marne-la-Vallée, 5, boulevard Descartes, Champs-sur-Marne, F-77454 Marne-la-Vallée Cedex 2.
e-mail address: dominique.perrin@esiee.fr

IRIF, CNRS and Université Paris-Diderot,, Case 7014, 75205 Paris Cedex 13, France.
e-mail address: jean-eric.pin@irif.fr

ABSTRACT. Difference hierarchies were originally introduced by Hausdorff and they play an important role in descriptive set theory. In this survey paper, we study difference hierarchies of regular languages. The first sections describe standard techniques on difference hierarchies, mostly due to Hausdorff. We illustrate these techniques by giving decidability results on the difference hierarchies based on shuffle ideals, strongly cyclic regular languages and the polynomial closure of group languages.

Dedicated to the memory of Zoltán Ésik.

1. INTRODUCTION

Difference hierarchies were originally introduced by Hausdorff [12, 13, 14]. They play an important role in descriptive set theory [24, Section 11] and also yield a hierarchy on complexity classes known as the Boolean hierarchy [15, Section 3], [26, Section 2], [3], [2, Section 3]. Difference hierarchies were also used in the study of ω-regular languages [4, 6, 8, 7, 9, 25].

The aim of this paper is to survey difference hierarchies of regular languages. Decidability questions for difference hierarchies over regular languages were studied in [10] and more recently by Glasser, Schmitz and Selivanov in [11]. This latter article is the reference paper on this topic and contains an extensive bibliography, to which we refer the interested reader. However, paper [11] focuses on forbidden patterns in automata, a rather different perspective than ours.

Received by the editors Tuesday 28th February, 2017 02:24.

1998 ACM Subject Classification: Formal languages and automata theory, Regular languages, Algebraic language theory.

Key words and phrases: Difference hierarchy, regular language, ordered syntactic monoid.

The third author is partially funded from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 670624). The first and third authors are partially funded by the DeLTA project (ANR-16-CE40-0007).

LOGICAL METHODS
IN COMPUTER SCIENCE

DOI:10.2168/LMCS-???

© O. Carton, D. Perrin, and J.-É. Pin
Creative Commons

Our paper is organised as follows. Prerequities are presented in Section 2. Section 3 covers the results of Hausdorff on difference hierarchies and Section 4 is a brief summary on closure operators. The results on approximation of Section 5, first presented in [5], lead in some cases to convenient algorithms to compute chain hierarchies. Section 6 on chains is inspired by results of descriptive set theory. Two results that are not addressed in [11] are presented in Sections 7 and 9. The first one, taken from [5], deals with cyclic and strongly cyclic regular languages. The second one studies the difference hierarchy of the polynomial closure of a lattice of regular languages and appears to be new.

2. PREREQUISITES

In this section, we briefly recall the following notions: upsets, ordered monoids, stamps and syntactic objects.

Let E be a preordered set. An *upset* of E is a subset U of E such that the conditions $x \in U$ and $x \leqslant y$ imply $y \in U$. An *ordered monoid* is a monoid M equipped with a partial order $\leqslant$ compatible with the product on M: for all $x, y, z \in M$, if $x \leqslant y$ then $zx \leqslant zy$ and $xz \leqslant yz$.

A *stamp* is a surjective monoid morphism $\varphi : A^* \to M$ from a finitely generated free monoid A^* onto a finite ordered monoid M. If M is an ordered monoid, φ is called an *ordered stamp*.

The *restricted direct product* of two stamps $\varphi_1 : A^* \to M_1$ and $\varphi_2 : A^* \to M_2$ is the stamp φ with domain A^* defined by $\varphi(a) = (\varphi_1(a), \varphi_2(a))$. The image of φ is an [ordered] submonoid of the [ordered] monoid $M_1 \times M_2$.



Stamps and ordered stamps are used to recognise languages. A language L of A^* is *recognised by a stamp* $\varphi : A^* \to M$ if there exists a subset P of M such that $L = \varphi^{-1}(P)$. It is *recognised by an ordered stamp* $\varphi : A^* \to M$ if there exists an upset U of M such that $L = \varphi^{-1}(U)$.

The syntactic preorder of a language was first introduced by Schützenberger in [22, p. 10]. Let L be a language of A^*. The *syntactic preorder* of L is the relation $\leqslant_L$ defined on A^* by $u \leqslant_L v$ if and only if, for every $x, y \in A^*$,

$$xuy \in L \implies xvy \in L.$$

The associated equivalence relation $\sim_L$, defined by $u \sim_L v$ if $u \leqslant_L v$ and $v \leqslant_L u$, is the *syntactic congruence* of L and the quotient monoid $M(L) = A^*/\sim_L$ is the *syntactic monoid* of L. The natural morphism $\eta : A^* \to A^*/\sim_L$ is the *syntactic stamp* of L. The *syntactic image* of L is the set $P = \eta(L)$.

The *syntactic order* $\leqslant_P$ is defined on $M(L)$ as follows: $u \leqslant_P v$ if and only if for all $x, y \in M$,

$$xuy \in P \implies xvy \in P$$

The partial order $\leqslant_P$ is stable and the resulting ordered monoid $(M, \leqslant_P)$ is called the *ordered syntactic monoid* of L. Note that P is now an upset of $(M, \leqslant_P)$ and η becomes an ordered stamp, called the ordered syntactic stamp of L.

3. Difference hierarchies

Let E be a set. In this article, a *lattice* is simply a collection of subsets of E containing $\emptyset$ and E and closed under taking finite unions and finite intersections. A *lattice* closed under complement is a *Boolean algebra*. Thorough this paper, we adopt Hausdorff's convention to denote union additively, set difference by a minus sign and intersection as a product.

Let $\mathcal{F}$ be a set of subsets of E containing the empty set. We set $\mathcal{B}_0(\mathcal{F}) = \{\emptyset\}$ and, for each integer $n \geqslant 1$, we let $\mathcal{B}_n(\mathcal{F})$ denote the class of all sets of the form

$$X = X_1 - X_2 + \cdots \pm X_n \tag{3.1}$$

where the sets X_i are in $\mathcal{F}$ and satisfy $X_1 \supseteq X_2 \supseteq X_3 \supseteq \cdots \supseteq X_n$. By convention, the expression on the right hand side of (3.1) should be evaluated from left to right, but given the conditions on the X_i's, it can also be evaluated as

$$(X_1 - X_2) + (X_3 - X_4) + (X_5 - X_6) + \cdots \tag{3.2}$$

Since the empty set belongs to $\mathcal{F}$, one has $\mathcal{B}_n(\mathcal{F}) \subseteq \mathcal{B}_{n+1}(\mathcal{F})$ for all $n \geqslant 0$ and the classes $\mathcal{B}_n(\mathcal{F})$ define a hierarchy within the Boolean closure of $\mathcal{F}$. Moreover, the following result, due to Hausdorff [13], holds:

Theorem 3.1. *Let $\mathcal{F}$ be a lattice of subsets of E. The union of the classes $\mathcal{B}_n(\mathcal{F})$ for $n \geqslant 0$ is the Boolean closure of $\mathcal{F}$.*

Proof. Let $\mathcal{B}(\mathcal{F}) = \cup_{n \geqslant 1} \mathcal{B}_n(\mathcal{F})$. We have $\mathcal{B}_1(\mathcal{F}) = \mathcal{F}$ and thus $\mathcal{F} \subseteq \mathcal{B}$. It is therefore enough to prove that $\mathcal{B}(\mathcal{F})$ is closed under complement and finite intersection. If $X = X_1 - X_2 + \cdots \pm X_n$, one has

$$E - X = E - X_1 + X_2 - \cdots \mp X_n$$

and thus $X \in \mathcal{B}(\mathcal{F})$ implies $E - X \in \mathcal{B}(\mathcal{F})$.

Let $X = X_1 - X_2 + \cdots \pm X_n$ and $Y = Y_1 - Y_2 + \cdots \pm Y_m$ be two elements of $\mathcal{B}$. Let

$$Z = Z_1 - Z_2 + \cdots \pm Z_{n+m}$$

with

$$Z_k = \sum_{\substack{i+j=k+1 \\ i \text{ and } j \text{ not both even}}} X_i Y_j$$

Therefore

$$\begin{aligned}
Z_1 &= X_1 Y_1, \\
Z_2 &= X_1 Y_2 + X_2 Y_1, \\
Z_3 &= X_1 Y_3 + X_3 Y_1, \\
Z_4 &= X_1 Y_4 + X_2 Y_3 + X_3 Y_2 + X_4 Y_1, \\
&\vdots \\
Z_{n+m-1} &= \begin{cases} X_n Y_m & \text{if } m \text{ and } n \text{ are not both even} \\ \emptyset & \text{otherwise} \end{cases}
\end{aligned}$$

We claim that $Z = XY$. To prove the claim, consider for each set $X = X_1 - X_2 + \cdots \pm X_n$ associated with the decreasing sequence $X_1, \ldots, X_n$ of subsets of E, the function μ_X defined on E by

$$\mu_X(x) = \max\{i \geqslant 1 \mid x \in X_i\}$$

with the convention that $\mu_X(x) = 0$ if $x \in E - X_1$. Then $x \in X$ if and only if $\mu_X(x)$ is odd. We now evaluate $\mu_Z(x)$ as a function of $i = \mu_X(x)$ and $j = \mu_Y(x)$. We first observe that if $k \geqslant i + j$, then $x \notin Z_k$. Next, if i and j are not both even, then $x \in X_iY_j$ and $X_iY_j \subseteq Z_{i+j-1}$, whence $\mu_Z(x) = i + j - 1$. Finally, if i and j are both even, then $x \notin Z_{i+j-1}$ and thus $\mu_Z(x)$ is either equal to 0 or to $i + j - 2$. Summarizing the various cases, we observe that $\mu_X(x)$ and $\mu_Y(x)$ are both odd if and only if $\mu_Z(s)$ is odd, which proves the claim. It follows that $\mathcal{B}(\mathcal{F})$ is closed under complement. □

An equivalent definition of $\mathcal{B}_n$ was given by Hausdorff [14]. Let $X \triangle Y$ denote the symmetric difference of two subsets X and Y of E.

Proposition 3.2. *For every $n \geqslant 0$, $\mathcal{B}_n(\mathcal{F}) = \{X_1 \triangle X_2 \triangle \cdots \triangle X_n \mid X_i \in \mathcal{F}\}$.*

Proof. Indeed, if $X = X_1 - X_2 + \cdots \pm X_n$ with $X_1 \supseteq X_2 \supseteq X_3 \supseteq \cdots \supseteq X_n$, then $X = X_1 \triangle X_2 \triangle \cdots \triangle X_n$. In the opposite direction, if $X = X_1 \triangle X_2 \triangle \cdots \triangle X_n$, then $X = Z_1 - Z_2 + \cdots \pm Z_n$ where $Z_k = \sum_{i_1, \ldots, i_k \text{ distincts}} X_{i_1} \cdots X_{i_k}$. □

4. Closure operators

We review in this section the definition and the basic properties of closure operators.

Let E be a set. A map $X \to \overline{X}$ from $\mathcal{P}(E)$ to itself is a *closure operator* if it is *extensive, idempotent and isotone*, that is, if the following properties hold for all $X, Y \subseteq E$:

(1) $X \subseteq \overline{X}$ (extensive)

(2) $\overline{\overline{X}} = \overline{X}$ (idempotent)

(3) $X \subseteq Y$ implies $\overline{X} \subseteq \overline{Y}$ (isotone)

A set $F \subseteq E$ is *closed* if $\overline{F} = F$. If F is closed, and if $X \subseteq F$, then $\overline{X} \subseteq \overline{F} = F$. It follows that $\overline{X}$ is the smallest closed set containing X. This justifies the terminology "closure". Actually, closure operators can be characterised by their closed sets.

Proposition 4.1. *A set of subsets of E is the set of closed sets for some closure operator on E if and only if it is closed under (possibly infinite) intersection.*

Proof. Let $X \to \overline{X}$ be a closure operator and let $(F_i)_{i \in I}$ be a family of closed subsets of E. Since a closure is isotone, $\overline{\bigcap_{i \in I} F_i} \subseteq \overline{F_i} = F_i$. It follows that $\overline{\bigcap_{i \in I} F_i} \subseteq \bigcap_{i \in I} F_i$ and thus $\bigcap_{i \in I} F_i$ is closed.

Conversely, given a set $\mathcal{F}$ of subsets of E closed under intersection, denote by $\overline{X}$ the intersection of all elements of $\mathcal{F}$ containing X. Then the map $X \to \overline{X}$ is a closure operator. □

In particular, $\overline{X \cap Y} \subseteq \overline{X} \cap \overline{Y}$, but the inclusion may be strict.

Example 4.1. The trivial closure is the application defined by

$$\overline{X} = \begin{cases} \emptyset & \text{if } X = \emptyset \\ E & \text{otherwise} \end{cases}$$

For this closure, the only closed sets are the empty set and E.

Example 4.2. If E is a topological space, the closure in the topological sense is a closure operator.

Example 4.3. The convex hull is a closure operator. However, it is not induced by any topology, since the union of two convex sets is not necessarily convex.

The *intersection* of two closure operators $X \to \overline{X}^1$ and $X \to \overline{X}^2$ is the function $X \to \overline{X}^3$ defined by $\overline{X}^3 = \overline{X}^1 \cap \overline{X}^2$.

Proposition 4.2. *The intersection of two closure operators is a closure operator.*

Proof. Let $\overline{}^3$ be the intersection of $\overline{}^1$ and $\overline{}^2$. First, since $X \subseteq \overline{X}^1$ and $X \subseteq \overline{X}^2$, one has $X \subseteq \overline{X}^3 = \overline{X}^1 \cap \overline{X}^2$. In particular, $\overline{X}^3 \subseteq \overline{\overline{X}^3}^3$. Secondly, since $\overline{X}^1 \cap \overline{X}^2 \subseteq \overline{X}^1$, $\overline{\overline{X}^1 \cap \overline{X}^2}^1 \subseteq \overline{\overline{X}^1}^1 = \overline{X}^1$. Similarly, $\overline{\overline{X}^1 \cap \overline{X}^2}^2 \subseteq \overline{X}^2$. It follows that

$$\overline{\overline{X}^3}^3 = \overline{\overline{X}^1 \cap \overline{X}^2}^1 \cap \overline{\overline{X}^1 \cap \overline{X}^2}^2 \subseteq \overline{X}^1 \cap \overline{X}^2 = \overline{X}^3$$

and hence $\overline{X}^3 = \overline{\overline{X}^3}^3$. Finally, if $X \subseteq Y$, then $\overline{X}^1 \subseteq \overline{Y}^1$ and $\overline{X}^2 \subseteq \overline{Y}^2$, and therefore $\overline{X}^3 \subseteq \overline{Y}^3$. □

Let us concluse this section by giving a few examples of closure operators occurring in the theory of formal languages.

Example 4.4. Iteration. The map $L \to L^*$ is a closure operator. Similarly, the map $L \to L^+$, where L^+ denotes the subsemigroup generated by L, is a closure operator.

Example 4.5. Shuffle ideal. The *shuffle product* (or simply *shuffle*) of two languages L_1 and L_2 over A is the language

$$L_1 \sqcup\!\sqcup L_2 = \{w \in A^* \mid w = u_1 v_1 \cdots u_n v_n \text{ for some words } u_1, \ldots, u_n \\ v_1, \ldots, v_n \text{ of } A^* \text{ such that } u_1 \cdots u_n \in L_1 \text{ and } v_1 \cdots v_n \in L_2\}.$$

The shuffle product defines a commutative and associative operation over the set of languages over A. Given a language L, the language $L \sqcup\!\sqcup A^*$ is called the *shuffle ideal* generated by L and it is easy to see that the map $L \to L \sqcup\!\sqcup A^*$ is a closure operator.

This closure operator can be extended to infinite words in two ways: the *finite and infinite shuffle ideals* generated by an ω-language X are respectively:

$$X \sqcup\!\sqcup A^* = \{y_0 x_1 y_1 \cdots x_n y_n x \mid x_1 \cdots x_n x \in X\}$$
$$X \sqcup\!\sqcup A^\omega = \{y_0 x_1 y_1 x_2 \cdots \mid x_1 x_2 \cdots \in X\}$$

The maps $X \to X \sqcup\!\sqcup A^*$ and $X \to X \sqcup\!\sqcup A^\omega$ are both closure operators.

Example 4.6. Ultimate closure. The *ultimate closure* of a language X of infinite words is defined by:

$$\mathrm{Ult}(X) = \{ux \mid u \in A^* \text{ and } vx \in X \text{ for some } v \in A^*\}$$

The map $X \to \mathrm{Ult}(X)$ is a closure operator.

5. APPROXIMATION

In this section, we consider a set $\mathcal{F}$ of closed sets of E containing the empty set. It follows that the corresponding closure operator satisfies the condition $\overline{\emptyset} = \emptyset$. We first define the notion of an *approximation* of a set by a chain of closed sets. Then the existence of a best approximation will be established.

Definition 5.1. Let L be a subset of E. A chain $F_1 \supseteq F_2 \supseteq \cdots \supseteq F_n$ of closed sets is an *n-approximation of L* if the following inclusions hold:

$$F_1 - F_2 \subseteq F_1 - F_2 + F_3 - F_4 \subseteq \cdots \subseteq F_1 - F_2 + \cdots + F_{2k-1} - F_{2k} \subseteq \cdots \subseteq L \subseteq \cdots \subseteq F_1 - F_2 + F_3 - \cdots + F_{2k+1} \subseteq \cdots \subseteq F_1 - F_2 + F_3 \subseteq F_1$$

There is a natural order among the n-approximations of a given set L. An n-approximation $F_1 \supseteq F_2 \supseteq \cdots \supseteq F_n$ of L is said to be *better* than an n-approximation $F'_1 \supseteq F'_2 \supseteq \cdots \supseteq F'_n$ if, for all k such that $2k + 1 \leqslant n$,

$$F_1 - F_2 + F_3 - \cdots + F_{2k+1} \subseteq F'_1 - F'_2 + F'_3 - \cdots + F'_{2k+1}$$

and

$$F'_1 - F'_2 + \cdots + F'_{2k-1} - F'_{2k} \subseteq F_1 - F_2 + \cdots + F_{2k-1} - F_{2k}$$

We will need the following elementary lemma:

Lemma 5.1. *Let X, Y and Z be subsets of E.*

(1) *The conditions $X - Y \subseteq Z$ and $X - Z \subseteq Y$ are equivalent,*
(2) *If $Y \subseteq X$ and $X - Y \subseteq Z$, then $X - Z = Y - Z$.*

The description of the best approximation of a subset L of E requires the introduction of two auxiliary functions. For every subset X of E, set

$$f(X) = \overline{X - L} \quad \text{and} \quad g(X) = \overline{X \cap L} \tag{5.1}$$

The key properties of these functions are formulated in the following lemma.

Lemma 5.2. *For every subset X of E, $X - f(X) \subseteq L$ and $X - g(X) \subseteq L^c$. Furthermore, the following properties hold:*

(1) *if $X \supseteq Y \supseteq L$, then $f(X) \supseteq f(Y)$ and $X - f(X) \subseteq Y - f(Y) \subseteq L$,*
(2) *if $X \subseteq Y \subseteq L$, then $g(X) \supseteq g(Y)$ and $X - g(X) \subseteq Y - g(Y) \subseteq L$.*

Proof. Since $g(X) = \overline{X - L^c}$, the results concerning g can be deduced from those concerning f by taking the complement. Let us prove the statements involving f. The first part of the lemma follows from a simple computation: $X - f(X) = X - \overline{X - L} \subseteq X - (X - L) = X \cap L \subseteq L$.

Suppose now that $X \supseteq Y \supseteq L$. Then $X - L \supseteq Y - L$ and thus $f(X) \supseteq f(Y)$. Furthermore, $X - Y \subseteq X - L \subseteq \overline{X - L} = f(X)$. Therefore, by Lemma 5.1, $X - f(X) = Y - f(X) \subseteq Y - f(Y)$. □

Lemma 5.3. *Let $F_1 \supseteq F_2 \supseteq \cdots \supseteq F_n$ be an n-approximation of L and, for $1 \leqslant k \leqslant n$, let $S_k = F_1 - F_2 + \cdots \pm F_k$. Then, for $1 \leqslant k \leqslant n$,*

$$\begin{cases} f(S_k) = f(F_k) \text{ if } k \text{ is odd} \\ g(S_k^c) = g(F_k) \text{ if } k \text{ is even} \end{cases} \tag{5.2}$$

Proof. If $k = 1$, then $S_1 = F_1$ and the result is trivial. Suppose that $k > 1$. If k is odd, $S_{k-1} \subseteq L$ and thus $S_k - L = (S_{k-1} + F_k) - L = F_k - L$. It follows that $f(S_k) = f(F_k)$. If k is even, $L \subseteq S_{k-1}$ and thus $S_k^c \cap L = (S_{k-1}^c + F_k) \cap L = F_k \cap L$. Therefore $g(S_k^c) = g(F_k)$. □

Define a sequence $(L_n)_{n\geqslant 0}$ of subsets of E by $L_0 = E$ and, for all $n \geqslant 0$,

$$L_{n+1} = \begin{cases} f(L_n) & \text{if } n \text{ is odd} \\ g(L_n) & \text{if } n \text{ is even} \end{cases} \tag{5.3}$$

The next theorem expresses the fact that the sequence $(L_n)_{n\geqslant 0}$ is the best approximation of L as a Boolean combination of closed sets. In particular, if $L_n = \emptyset$ for some $n > 0$, then $L \in \mathcal{B}_{n-1}(\mathcal{F})$.

Theorem 5.4. *For every $n > 0$, the sequence $(L_k)_{1\leqslant k\leqslant n}$ is the best n-approximation of L.*

Proof. We first show that the sequence $(L_k)_{1\leqslant k\leqslant n}$ is an n-approximation of L. Set, for $k > 0$, $S_k = L_1 - L_2 + \cdots \pm L_k$. First, every L_k is closed by construction. We show that $L_{k+1} \subseteq L_k$ by induction on k. This is true for $k = 0$ since $L_0 = E$. Now, if k is even, $L_{k+1} = \overline{L_k \cap L} \subseteq \overline{L_k} = L_k$ and if k is odd, $L_{k+1} = \overline{L_k - L} \subseteq \overline{L_k} = L_k$.

Next, by Lemma 5.2, the relations $L_{2k-1} - L_{2k} = L_{2k-1} - f(L_{2k-1}) \subseteq L$ hold for every $k > 0$, and similarly, $L_{2k} - L_{2k+1} = L_{2k} - g(L_{2k}) \subseteq L^c$. It follows that $S_{2k} \subseteq L$. Furthermore $S^c_{2k+1} = (L_0 - L_1) + (L_2 - L_3) + \cdots + (L_{2k} - L_{2k+1}) \subseteq L^c$ and thus $L \subseteq S_{2k+1}$.

We now show that the sequence $(L_k)_{1\leqslant k\leqslant n}$ is the best approximation of L. Let $(L'_k)_{1\leqslant k\leqslant n}$ be another n-approximation of L. Set, for $k > 0$, $S'_k = L'_1 - L'_2 + \cdots \pm L'_k$. Then, by definition, $L \subseteq L'_1$ and thus

$$S_1 = L_1 = \overline{L} \subseteq \overline{L'_1} = L'_1 = S'_1.$$

Let k be an even number and suppose by induction that $S_{k-1} \subseteq S'_{k-1}$. Then, by definition of an approximation, $S'_k = S'_{k-1} - L'_k \subseteq L$, and thus by Lemma 5.1, $S'_{k-1} - L \subseteq L'_k$. It follows $f(S'_{k-1}) = \overline{S'_{k-1} - L} \subseteq \overline{L'_k} = L'_k$. Therefore, by Lemma 5.2,

$$S'_k = S'_{k-1} - L'_k \subseteq S'_{k-1} - f(S'_{k-1}) \subseteq S_{k-1} - f(S_{k-1}).$$

Now, by Lemma 5.3, $f(S_{k-1}) = f(L_{k-1}) = L_k$, whence $S'_k \subseteq S_{k-1} - L_k = S_k$.

Similarly, $L \subseteq S'_{k+1} = S'_k + L'_{k+1}$ and thus $S'^c_k - L^c \subseteq L'_{k+1}$. It follows that

$$g(S'^c_k) = \overline{S'^c_k - L^c} \subseteq \overline{L'_{k+1}} = L'_{k+1}.$$

Therefore, one gets by Lemma 5.1,

$$S'^c_{k+1} = S'^c_k - L'_{k+1} \subseteq S'^c_k - g(S'^c_k) \subseteq S^c_k - g(S^c_k).$$

Now, by Lemma 5.3, $g(S^c_k) = g(L_k) = L_{k+1}$. Thus $S'^c_{k+1} \subseteq S^c_k - L_{k+1} = S^c_{k+1}$, whence $S_{k+1} \subseteq S'_{k+1}$. □

Corollary 5.5. *Let $\mathcal{F}$ be a set of subsets of E closed under (possibly infinite) intersection and containing the empty set. Let $(L_k)_{1\leqslant k\leqslant n}$ be the best n-approximation of L with respect to $\mathcal{F}$. Then $L \in \mathcal{B}_{n-1}(\mathcal{F})$ if and only if $L_n = \emptyset$ and in this case*

$$L = L_1 - L_2 + \cdots \pm L_{n-1} \tag{5.4}$$

Proof. If $L \in \mathcal{B}_{n-1}(\mathcal{F})$, then $L = F_1 - F_2 + \cdots \pm F_{n-1}$ with $F_1, \ldots, F_{n-1} \in \mathcal{F}$. Let $F_n = \emptyset$. Then the sequence $(F_k)_{1\leqslant k\leqslant n}$ is an n-approximation of L. Since $(L_k)_{1\leqslant k\leqslant n}$ is the best n-approximation of L, one has $L = L_1 - L_2 + \cdots \pm L_{n-1}$. Thus, with the notation of Lemma 5.3,

$$\begin{cases} f(L_{n-1}) = f(L) = \emptyset \text{ if } n-1 \text{ is odd} \\ g(L_{n-1}) = g(L^c) = \emptyset \text{ if } n-1 \text{ is even} \end{cases} \tag{5.5}$$

Therefore, $L_n = \emptyset$ by (5.3).

Conversely, suppose that $L_n = \emptyset$. If $n = 2k$, then

$$L = (L_1 - L_2) + \cdots + (L_{2k-1} - L_{2k}) \subseteq L \subseteq (L_1 - L_2) + \cdots + (L_{2k-3} - L_{2k-2}) + L_{2k-1}$$

If $n = 2k + 1$, then

$$L = (L_1 - L_2) + \cdots + (L_{2k-1} - L_{2k}) \subseteq L \subseteq (L_1 - L_2) + \cdots + (L_{2k-1} - L_{2k}) + L_{2k+1}$$

In both cases, one gets $L = L_1 - L_2 + \cdots \pm L_{n-1}$ and thus $L \in \mathcal{B}_{n-1}(\mathcal{F})$. □

Let us illustrate this corollary by a concrete example.

Example 5.1. Let $A = \{a, b, c\}$ and let $\mathcal{L}$ be the lattice of shuffle ideals. If L is the language $\{1, a, b, c, ab, bc, abc\}$, a straightforward computation gives

$$L_0 = A^*$$
$$L_1 = g(L_0) = A^* \sqcup (L_0 \cap L) = A^* \sqcup L = A^*$$
$$L_2 = f(L_1) = A^* \sqcup (L_1 - L) = A^* \sqcup \{aa, ac, ba, bb, ca, cb, cc\}$$
$$L_3 = g(L_2) = A^* \sqcup (L_2 \cap L) = A^* \sqcup abc$$
$$L_4 = f(L_3) = A^* \sqcup (L_3 - L) = A^* \sqcup \{aabc, abac, abca, babc, abbc, abcb, cabc, acbc, abcc\}$$
$$L_5 = g(L_4) = A^* \sqcup (L_4 \cap L) = \emptyset$$

It follows that $L = L_1 - L_2 + L_3 - L_4$ and $L \in \mathcal{B}_4(\mathcal{L})$, but $L \notin \mathcal{B}_3(\mathcal{L})$.

It is also possible to use the approximation algorithm for a set $\mathcal{L}$ of subsets of E closed under (possibly infinite) union and containing the set E. In this case, the set

$$\mathcal{L}^c = \{L^c \mid L \in \mathcal{L}\}$$

is closed under (possibly infinite) intersection and contains the empty set. Consequently, the approximation algorithm can be applied to $\mathcal{L}^c$ but it describes the difference hierarchy $\mathcal{B}_n(\mathcal{L}^c)$. To recover the difference hierarchy $\mathcal{B}_n(\mathcal{L})$, the following algorithm can be used. First compute the best $\mathcal{L}^c$-approximation of even length of L and the best $\mathcal{L}^c$-approximation of odd length of L^c, say

$$L = L_1^c - L_2^c + \cdots \pm L_n^c \tag{5.6}$$
$$L^c = F_1^c - F_2^c + \cdots \pm F_m^c \tag{5.7}$$

with n even, m odd, $L_i, F_i \in \mathcal{L}$ and L_n and L_m possibly empty to fill the parity requirements. Now L admits the following $\mathcal{L}$-decompositions, where L_1 and F_1 are possibly empty (and consequently deleted):

$$L = L_n - L_{n-1} + \cdots \pm L_1 \tag{5.8}$$
$$= F_m - F_{m-1} + \cdots \pm F_1 \tag{5.9}$$

It remains to take the shortest of the two expressions to get the best $\mathcal{L}$-approximation of L.

6. Chains and difference hierarchies

Chains can be defined on any ordered set. We first give their definition, then establish a connection with the difference hierarchy.

Definition 6.1. Let $(E, \leqslant)$ be a partially ordered set and let X be a subset of E. A *chain* of E is a strictly increasing sequence

$$x_0 < x_1 < \ldots < x_{m-1}$$

of elements of E. It is called an *X-chain* if x_0 is in X and the x_i's are alternatively elements of X and of its complement X^c. The integer m is called the *length* of the chain. We let $m(X)$ denote the maximal length of an X-chain.

There is a subtle connection between chains and difference hierarchies of regular languages. Let M be a finite ordered monoid and let $\varphi : A^* \to M$ be a surjective monoid morphism. Let

$$\mathcal{L} = \{\varphi^{-1}(U) \mid U \text{ is an upset of } M\}$$

By definition, every language of $\mathcal{L}$ is recognised by the ordered monoid M.

Theorem 6.1. *If there exists a subset P of M such that $L = \varphi^{-1}(P)$ and $m(P) \leqslant n$, then L belongs to $\mathcal{B}_n(\mathcal{L})$.*

Before starting the proof, let us clarify a delicate point. The condition $L = \varphi^{-1}(P)$ means that L is recognised by the *monoid* M. It does not mean that L is recognised by the *ordered monoid* M, a property which would require P to be an upset.

Proof. For each $s \in M$, let $m(P, s)$ be the maximal length of a P-chain ending with s. Finally, let, for each $k > 0$,

$$U_k = \{s \in M \mid m(P, s) \geqslant k\}$$

We claim that U_k is an upset of M. Indeed, if $s \in U_k$, there exists a P-chain $x_0 < x_1 < \cdots < x_{r-1} = s$ of length $r \geqslant k$. Let t be an element of M such that $s \leqslant t$. If s and t are not simultaneously in P, then $x_0 < x_1 < \cdots < x_{r-1} < t$ is a P-chain of length $r + 1 \geqslant k$. Otherwise, $x_0 < x_1 < \cdots < x_{r-2} < t$ is a P-chain of length $r \geqslant k$. Thus $m(P, t) \geqslant k$, and $t \in U_k$, proving the claim.

We now show that

$$P = U_1 - U_2 + U_3 - U_4 \cdots \pm U_n \tag{6.1}$$

First observe that $s \in P$ if and only if $m(P, s)$ is odd. Since $m(P) \leqslant n$, one has $m(P, s) \leqslant n$ for every $s \in M$ and thus $U_{n+1} = \emptyset$. Formula (6.1) follows, since for each $r \geqslant 0$,

$$\{s \in M \mid m(P, s) = r\} = U_r - U_{r+1}.$$

Let, for $1 \leqslant i \leqslant n$, $L_i = \varphi^{-1}(U_i)$. Since U_i is an upset, each L_i belongs to $\mathcal{L}$. Moreover, one gets from (6.1) the formula

$$L = L_1 - L_2 + L_3 \cdots \pm L_n \tag{6.2}$$

which shows that $L \in \mathcal{B}_n(\mathcal{L})$. □

We now establish a partial converse to Theorem 6.1. A *lattice of regular languages* is a set $\mathcal{L}$ of regular languages of A^* containing $\emptyset$ and A^* and closed under finite union and finite intersection. Let $\mathcal{L}$ be a lattice of regular languages of A^*.

Theorem 6.2. *If a language L belongs to $\mathcal{B}_n(\mathcal{L})$, then there exist an ordered stamp $\eta : A^* \to M$ and a subset P of M satisfying the following conditiosn:*

(1) *φ is a restricted product of syntactic ordered stamps of members of $\mathcal{L}$,*
(2) *$L = \eta^{-1}(P)$,*
(3) *$m(P) \leqslant n$.*

Proof. If $L \in \mathcal{B}_n(\mathcal{L})$, then

$$L = L_1 - L_2 + L_3 \cdots \pm L_n$$

with $L_1 \supseteq L_2 \supseteq \cdots \supseteq L_n$ and $L_i \in \mathcal{L}$. Let $\eta_i : A^* \to (M_i, \leqslant_i)$ be the syntactic morphism of L_i and let $U_i = \eta_i(L_i)$. Then each U_i is an upset of M_i and $L_i = \eta_i^{-1}(U_i)$. Let $\eta : A^* \to M$ be the restricted product of the stamps η_i. Condition (1) is satisfied by construction.

Observe that if $\eta(u) = (s_1, \ldots, s_n)$ is an element of M, the condition $s_{i+1} \in P_{i+1}$ is equivalent with $u \in L_{i+1}$, and since L_{i+1} is a subset of L_i, this condition also implies $u \in L_i$ and $s_i \in P_i$. Consequently, for each element $s = (s_1, \ldots, s_n)$ of M, there exists a unique $k \in \{0, \ldots, n\}$ such that

$$s_1 \in P_1, \ldots, s_k \in P_k, s_{k+1} \notin P_{k+1}, \ldots, s_n \notin P_n$$

This unique k is called the *cut* of s. Setting

$$P = \{s \in M \mid \text{the cut of } s \text{ is odd}\}$$

one gets

$$\eta^{-1}(P) = \bigcup_{k \text{ odd}} \Big((L_1 \cap \cdots \cap L_k) - L_{k+1} \Big) = \bigcup_{k \text{ odd}} (L_k - L_{k+1}) = L \tag{6.3}$$

which proves (2).

Let now $x_0 < x_1 < \cdots < x_{m-1}$ be a P-chain. Let, for $0 \leqslant i \leqslant m-1$, $x_i = (s_{i,1}, \ldots, s_{i,n})$ and let k_i be the cut of x_i. We claim that $k_{i+1} > k_i$. Indeed, since $x_i < x_{i+1}$, $s_{i,k_i} \leqslant_i s_{i+1,k_i}$ and since P_i is an upset, $s_{i,k_i} \in P_i$ implies $s_{i+1,k_i} \in P_{i+1}$, which proves that $k_{i+1} \geqslant k_i$. But since x_i and x_{i+1} are not simultaneously in P, their cuts must be different, which proves the claim. Since $x_0 \in P$, the cut of x_0 is odd, and in particular, non-zero. It follows that $0 < k_0 < k_1 < \cdots < k_{m-1}$ and since the cuts are numbers between 0 and n, $m \leqslant n$, which proves (3). □

It is tempting to try to improve Theorem 6.2 by taking for M the syntactic morphism of L and for φ the syntactic morphism of L. However, Example 5.1 ruins this hope. Indeed, let $F = \{1, a, b, c, ab, bc, abc\}$ be the set of factors of the word abc. Then the syntactic monoid of L can be defined as the set $F \cup \{0\}$ equipped with the product defined by

$$xy = \begin{cases} xy & \text{if } x, y \text{ and } xy \text{ are all in } F \\ 0 & \text{otherwise} \end{cases}$$

Now the syntactic image of L is equal to F. It follows that $M - F = \{0\}$ and thus, whatever order is taken on M, the length of a chain is bounded by 3, but in spite of that, L does not belong to $\mathcal{B}_3(\mathcal{L})$.

Therefore, if L is a regular language, the maximal length of an L-chain cannot be in general computed in the syntactic monoid of L. It follows that decidability questions on $\mathcal{B}_n(\mathcal{L})$, as presented in Section 8 below, cannot in general be solved just by inspecting the syntactic monoid. An exceptional case where the syntactic monoid suffices is presented in the next sections.

7. Cyclic and strongly cyclic regular languages

Cyclic and strongly cyclic regular languages are two classes of regular languages related to symbolic dynamic and first studied in [1]. It was shown in [5] that an appropriate notion of chains suffices to characterise the difference hierarchy based on the class of strongly cyclic regular languages. This contrasts with Section 6, in which the general results on chain did not lead to a full characterisation of difference hierarchies.

Let $\mathcal{A} = (Q, A, \cdot)$ be a finite (possibly incomplete) deterministic automaton. A word u *stabilises* a subset P of Q if $P \cdot u = P$. Given a subset P of Q, let $\mathrm{Stab}(P)$ be the set of all words

that stabilise P. The language $\mathrm{Stab}(\mathcal{A})$ that stabilises $\mathcal{A}$ is by definition the set of all words which stabilise at least one nonempty subset of Q.

Definition 7.1. A language is *strongly cyclic* if it stabilises some finite deterministic automaton.

Example 7.1. If $\mathcal{A}$ is the automaton represented in Figure 1, then

$$\mathrm{Stab}(\{1\}) = (b + aa)^*,\ \mathrm{Stab}(\{2\}) = (ab^*a)^*,\ \mathrm{Stab}(\{1,2\}) = a^*$$

and $\mathrm{Stab}(\mathcal{A}) = (b + aa)^* + (ab^*a)^* + a^*$.



Figure 1: The automaton $\mathcal{A}$.

One can show that the set of strongly cyclic languages of A^* form a lattice of languages but are not closed under quotients. For instance, as shown in Example 7.1, the language $L = (b + aa)^* + (ab^*a)^* + a^*$ is strongly cyclic, but its quotient $b^{-1}L = (b + aa)^*$ is not strongly cyclic, since $aa \in (b + aa)^*$ but $a \notin (b + aa)^*$.

We will also need the following characterisation [1, Proposition 7]:

Proposition 7.1. *Let $\mathcal{A} = (Q, A, E)$ be a deterministic automaton. A word u belongs to $\mathrm{Stab}(\mathcal{A})$ if and only if there is some state q of $\mathcal{A}$ such that for every integer n, the transition $q \cdot u^n$ exists.*

Strongly cyclic languages admit the following syntactic characterisation [1, Theorem 8]:

Proposition 7.2. *Let L be a nonfull regular language. The following conditions are equivalent:*

(1) *L is strongly cyclic,*

(2) *there is a morphism φ from A^* onto a finite monoid M with zero such that*

$$L = \varphi^{-1}(\{s \in M \mid s^\omega \neq 0\}),$$

(3) *the syntactic monoid M of L has a zero and its syntactic image is the set of all elements $s \in M$ such that $s^\omega \neq 0$.*

Proposition 7.2 leads to a simple syntactic characterisation of strongly cyclic languages. Recall that a language of A^* is *nondense* if there exists a word $u \in A^*$ such that $L \cap A^*uA^* = \emptyset$.

Proposition 7.3. *Let L be a regular language, let M be its syntactic monoid and let P be its syntactic image. Then L is strongly cyclic if and only if it satisfies the following conditions, for all $u, x, v \in M$:*

(S_1) *$ux^\omega v \in P$ implies $x^\omega \in P$,*

(S_2) *$x^\omega \in P$ if and only if $x \in P$.*

Furthermore, if these conditions are satisfied and if L is not the full language, then L is nondense.

Proof. Let L be a strongly cyclic language, let M be its syntactic monoid and let P be its syntactic image. If L is the full language, then the conditions (S_1) and (S_2) are trivially satisfied. If L is not the full language, then Proposition 7.2 shows that M has a zero and that $P = \{s \in M \mid s^\omega \neq 0\}$. Observing that $x^\omega = (x^\omega)^\omega$, one gets

$$x \in P \iff x^\omega \neq 0 \iff (x^\omega)^\omega \neq 0 \iff x^\omega \in P$$

which proves (S_2). Similarly, one gets

$$ux^\omega v \in P \iff (ux^\omega v)^\omega \neq 0 \implies x^\omega \neq 0 \iff x \in P$$

which proves (S_1).

Conversely, suppose that L satisfies (S_1) and (S_2). If L is full, then L is strongly cyclic. Otherwise, let $z \notin P$. Then $z^\omega \notin P$ by (S_1) and $uz^\omega v \notin P$ for all $u, v \in M$ by (S_2). This means that z is a zero of M and that $0 \notin P$. By Proposition 7.2, it remains to prove that $x \in P$ if and only if $x^\omega \neq 0$. First, if $x \in P$, then $x^\omega \in P$ by (S_2) and since $0 \notin P$, one has $x^\omega \neq 0$. Conversely, if $x^\omega \neq 0$, then $ux^\omega v \in P$ for some $u, v \in M$, since x^ω is not equivalent to 0 in the syntactic congruence of P. It follows that $x^\omega \in P$ by (S_1) and $x \in P$ by (S_2). □

We turn now to cyclic languages.

Definition 7.2. A subset of a monoid is said to be *cyclic* if it is closed under conjugation, power and root. Thus a subset P of a monoid M is cyclic if it satisfies the following conditions, for all $u, v \in M$ and $n > 0$:

(C_1) $u^n \in P$ if and only if $u \in P$,

(C_2) $uv \in P$ if and only if $vu \in P$.

This definition applies in particular to the case of a language of A^*.

Example 7.2. If $A = \{a, b\}$, the language b^* and its complement A^*aA^* are cyclic.

One can show that regular cyclic languages are closed under inverses of morphisms and under Boolean operations but not under quotients. For instance, the language $L = \{abc, bca, cab\}$ is cyclic, but its quotient $a^{-1}L = \{bc\}$ is not cyclic. Thus regular cyclic languages do not form a variety of languages. However, they admit the following straightforward characterisation in terms of monoids.

Proposition 7.4. *Let L be a regular language of A^*, let φ be a surjective morphism from A^* to a finite monoid M recognising L and let $P = \varphi(L)$. Then L is cyclic if and only if P is cyclic.*

Corollary 7.5. *Every strongly cyclic language is cyclic.*

Proof. Let L be a strongly cyclic language, let M be its syntactic monoid and let P be its syntactic image. By Proposition 7.3, P satisfies (S_1) and (S_2). It suffices now to prove that it satisfies (C_2). The sequence of equivalences

$$xy \in P \overset{(S_2)}{\iff} (xy)^\omega \in P \iff (xy)^\omega (xy)^\omega \in P \iff (xy)^{\omega-1}xy(xy)^{\omega-1}xy \in P$$
$$\iff ((xy)^{\omega-1}x)(yx)^\omega y \in P \overset{(S_1)}{\implies} (yx)^\omega \in P \overset{(S_2)}{\iff} yx \in P.$$

shows that $xy \in P$ implies $yx \in P$ and the opposite implication follows by symmetry. □

Proposition 7.2 implies that every strongly cyclic language is cyclic. Actually, for any regular cyclic language, there is a smallest strongly cyclic language containing it.

Proposition 7.6. *Let L be a regular cyclic language of A^*, let $\eta : A^* \to M$ be its syntactic stamp and let $P = \eta(L)$. There M has a zero and the language*

$$\overline{L} = \begin{cases} \eta^{-1}(\{s \mid s^\omega \neq 0\}) & \text{if } 0 \notin P, \\ A^* & \text{otherwise.} \end{cases}$$

is the smallest strongly cyclic language containing L.

Proof. If $0 \notin P$, then the language $\overline{L}$ is strongly cyclic by Proposition 7.2. Morevover, since L is cyclic, P is cyclic by Proposition 7.4. It follows that if $s \in P$, then $s^\omega \in P$ and in particular $s^\omega \neq 0$. Consequently, $\overline{L}$ contains L.

It remains to prove that $\overline{L}$ is the smallest strongly cyclic language containing L. Let X be a strongly cyclic language containing L and let u be a word of $\overline{L}$. Let $\mathcal{A} = (Q, A, E)$ be a deterministic automaton such that $X = \mathrm{Stab}(\mathcal{A})$. Setting $s = \eta(u)$, one has $s^\omega \neq 0$ by definition of $\overline{L}$. Consequently, $\eta(s)^n \neq 0$ for every integer n and there are two words x_n and y_n such that $x_n u^n y_n$ belongs to L. By Proposition 7.1, there is a state q_n of $\mathcal{A}$ such that the transition $q_n \cdot x_n u^n y_n$ is defined. The transition $(q_n \cdot x_n) \cdot u^n$ is thus defined for every n and by Proposition 7.1 again, the word u belongs to X. Thus $\overline{L} \subseteq X$ as required.

Suppose now that $0 \in P$ and let z be a word of L such that $\eta(z) = 0$. Let X be a strongly cyclic language containing L. If X is not full, then X is nondense by Proposition 7.3 and there exists a word $u \in A^*$ such that $A^*uA^* \cap X = \emptyset$. Since X contains L, one also gets $A^*uA^* \cap L = \emptyset$ and in particular $zu \notin L$. But this yieds a contradiction, since $\eta(zu) = \eta(z)\eta(u) = 0 \in P$ and thus $zu \in \eta^{-1}(P) = L$. Thus the only strongly cyclic language containing L is A^*. □

Given a finite monoid M, the Green's preorder relation $\leqslant_{\mathcal{J}}$ defined on M by

$$s \leqslant_{\mathcal{J}} t \text{ if and only if } s \in MtM, \text{ or equivalently, if there exists } u, v \in M \text{ such that } s = utv$$

is a preorder on M. The associated equivalence relation $\mathcal{J}$ is defined by

$$s \mathcal{J} t \text{ if } s \leqslant_{\mathcal{J}} t \text{ and } t \leqslant_{\mathcal{J}} s, \text{ or equivalently, if } MsM = MtM.$$

Corollary 7.7. *Let L be a regular cyclic language of A^*, let $\eta : A^* \to M$ be its syntactic stamp and let $P = \eta(L)$. Then L is strongly cyclic if and only if for all idempotent e, f of M, the conditions $e \in P$ and $e \leqslant_{\mathcal{J}} f$ imply $f \in P$.*

Proof. Suppose that L is strongly cyclic and let e, f be two idempotents of M such that $e \in P$ and $e \leqslant_{\mathcal{J}} f$. Let $u, v \in M$ be such that $e = ufv$. Since $f^\omega = f$, one gets $uf^\omega v \in P$ and thus $f \in P$ by Condition (S_1) of Proposition 7.3.

In the opposite direction, suppose that for all idempotents e, f of M, the conditions $e \in P$ and $e \leqslant_{\mathcal{J}} f$ imply $f \in P$. Since L is cyclic, it satisfies (C_1) and hence (S_2). We claim that it also satisfies (S_1). Indeed, $ux^\omega v \in P$ implies $(ux^\omega v)^\omega \in P$ by (S_2). Furthermore, since $(ux^\omega v)^\omega \leqslant_{\mathcal{J}} x^\omega$, one also has $x^\omega \in P$, and finally $x \in P$ by (S_2), which proves the claim. □

The precise connection between cyclic and strongly cyclic languages was given in [1].

Theorem 7.8. *A regular language is cyclic if and only if it is a Boolean combination of regular strongly cyclic languages.*

Theorem 7.8 motivates a detailed analysis of the difference hierarchy of the class $\mathcal{S}$ of strongly cyclic languages. This study relies on a careful analysis of the chains on the set of idempotents of a finite monoid, pre-ordered by the relation $\leqslant_{\mathcal{J}}$.

Let L be a regular language L of A^*, let φ be a surjective morphism from A^* onto a finite monoid M recognising L and let $P = \varphi(L)$.

Definition 7.3. A *P-chain of idempotents* is a sequence $(e_0, e_1, \ldots, e_{m-1})$ of idempotents of M such that

$$e_0 \leqslant_{\mathcal{J}} e_1 \leqslant_{\mathcal{J}} \cdots \leqslant_{\mathcal{J}} e_{m-1}$$

$e_0 \in P$ and, for $0 < i < m$, $e_i \in P$ if and only if $e_{i-1} \notin P$. The integer m is the length of the P-chain of idempotents.

We let $\ell(M, P)$ denote the maximal length of a P-chain of idempotents of M. We now prove that $\ell(M, P)$ just depends on the language recognised by M.

Theorem 7.9. *Let L be a regular language. Let $\varphi : A^* \to M$ and $\psi : A^* \to N$ be two stamps recognising L. If $P = \varphi(L)$ and $Q = \psi(L)$, then $\ell(M, P) = \ell(N, Q)$.*

Proof. It is sufficient to prove the result when φ is the syntactic stamp of L. Since the morphism ψ is surjective, M is a quotient of N and there is a surjective morphism $\pi : N \to M$ such that $\pi \circ \psi = \varphi$. It follows that

$$\pi(Q) = P \text{ and } \pi^{-1}(P) = Q. \tag{7.1}$$

We show that to any P-chain of idempotents in N, one can associate a Q-chain of idempotents of the same length in M and vice-versa.

Let $(e_0, \ldots, e_{m-1})$ be a Q-chain of idempotents in N and let $f_i = \pi(e_i)$ for $0 \leqslant i \leqslant m-1$. Since every monoid morphism preserves $\leqslant_{\mathcal{J}}$, the relations (7.1) show that $(f_0, \ldots, f_{m-1})$ is a P-chain of idempotents in M.

Let now $(f_0, \ldots, f_{m-1})$ be a P-chain of idempotents in M. Since $f_{i-1} \leqslant_{\mathcal{J}} f_i$, there exist for $1 \leqslant i \leqslant m-1$ elements u_i, v_i of M such that $u_i f_i v_i = f_{i-1}$. Let us choose an idempotent e_{m-1} such that $\pi(e_{m-1}) = f_{m-1}$ and some elements s_i and t_i of N such that $\pi(s_i) = u_i$ and $\pi(t_i) = v_i$. We now define a sequence of idempotents $(e_0, \ldots, e_{m-1})$ of N by setting

$$e_{m-2} = (s_{m-1}e_{m-1}t_{m-1})^\omega \qquad e_{m-3} = (s_{m-2}e_{m-2}t_{m-2})^\omega \qquad \cdots \qquad e_0 = (s_1e_1t_1)^\omega$$

By construction, $e_0 \leqslant_{\mathcal{J}} \cdots \leqslant_{\mathcal{J}} e_{m-1}$ and a straightforward induction shows that $\pi(e_i) = f_i$ for $0 \leqslant i \leqslant m-1$. Moreover the equalities (7.1) show that $e_i \in Q$ if and only if $f_i \in P$. It follows that $(e_0, \ldots, e_{m-1})$ is a Q-chain of idempotents of N and thus $\ell(M, P) = \ell(N, Q)$. □

Since the integers $\ell(M, P)$ only depend on L and not on the choice of the recognising monoid, let us define $\ell(L)$ as $\ell(M, P)$ where M $[P]$ is the syntactic monoid [image] of L. Note that by Corollary 7.7, a cyclic language L is strongly cyclic if and only if $\ell(L) = 1$. This is a special case of the following stronger result.

Theorem 7.10. *Let L be a regular cyclic language. Then $L \in \mathcal{B}_n(\mathcal{S})$ if and only if $\ell(L) \leqslant n$.*

We first prove the following lemma which states that the function ℓ is subadditive with respect to the symmetric difference.

Lemma 7.11. *If X and Y are regular languages, then $\ell(X \triangle Y) \leqslant \ell(X) + \ell(Y)$.*

Proof. Suppose that the languages X and Y are respectively recognised by the stamps $\varphi : A^* \to M$ and $\psi : A^* \to N$. Let P and Q be the images of X and Y in M and N, so that $X = \varphi^{-1}(P)$ and $Y = \psi^{-1}(Q)$. The language $X \triangle Y$ is recognised by the restricted product of the stamps φ and ψ, say $\gamma : A^* \to R$, and the image of $X \triangle Y$ in R is

$$T = R \cap \Big(P \times (N - Q) + (M - P) \times Q\Big).$$

Let $((e_0, f_0), \ldots, (e_{m-1}, f_{m-1}))$ be a T-chain of idempotents in R. Let us consider the set I $[J]$ of integers i for which exactly one of the idempotents e_{i-1} or e_i $[f_{i-1}$ or $f_i]$ belongs to P $[Q]$. Formally, we define the sets of integers I and J to be

$$I = \{1 \leqslant i \leqslant m-1 \mid e_{i-1} \in P \iff e_i \notin P\}$$
$$J = \{1 \leqslant i \leqslant m-1 \mid f_{i-1} \in Q \iff f_i \notin Q\}$$

Since the sequence $((e_0, f_0), \ldots, (e_{m-1}, f_{m-1}))$ is a T-chain in R, one has $e_0 \leqslant_{\mathcal{J}} \ldots \leqslant_{\mathcal{J}} e_{m-1}$ and $f_0 \leqslant_{\mathcal{J}} \ldots \leqslant_{\mathcal{J}} f_{m-1}$. Moreover, every integer i between 1 and $m-1$ belongs to exactly

one of the sets I or J. Otherwise, the idempotents (e_{i-1}, f_{i-1}) and (e_i, f_i) of R would be either both in T or both out of T. Let $I = \{i_1, \ldots, i_p\}$ and $J = \{j_1, \ldots, j_q\}$ with $i_1 < \cdots < i_p$ and $j_1 < \cdots < j_q$. Then $p + q = m - 1$.

Since $(e_0, f_0) \in T$, the conditions $e_0 \in P$ and $f_0 \notin Q$ are equivalent. By symmetry, suppose that $e_0 \in P$. Then $f_0 \notin Q$ and thus $f_1 \in Q$. Furthermore, the definitions of I and J give

$$\begin{array}{cccccccccc} e_0 \in P, & e_1 \in P, & \ldots & e_{i_1-1} \in P, & e_{i_1} \notin P, & \ldots & e_{i_2-1} \notin P, & e_{i_2} \in P, & \ldots \\ f_0 \notin P, & f_1 \notin P, & \ldots & f_{j_1-1} \notin P, & f_{j_1} \in P, & \ldots & f_{j_2-1} \in P, & f_{j_2} \notin P, & \ldots \end{array}$$

Then the sequence $(e_0, e_{i_1}, \ldots, e_{i_p})$ is a P-chain of idempotents in M and $(f_{j_1}, \ldots, f_q)$ is a Q-chain of idempotents in N. Therefore $p + 1 \leqslant \ell(X)$, $q \leqslant \ell(Y)$ and $m = p + 1 + q \leqslant \ell(X) + \ell(Y)$. Thus $\ell(X \triangle Y) \leqslant \ell(X) + \ell(Y)$. □

We can now complete the proof of Theorem 7.10. As usual, let $\eta : A^* \to M$ be the syntactic stamp of L and let $P = \eta(L)$. Let also $E(M)$ be the set of idempotents of M.

Proof. If $L \in \mathcal{B}_n(\mathcal{F})$, then $L = L_1 \triangle \cdots \triangle L_n$ for some strongly cyclic languages L_i. By Corollary 7.7, one has $\ell(L_i) = 1$ for $1 \leqslant i \leqslant n$ and thus $\ell(L) \leqslant n$ by Lemma 7.11.

Suppose now that $\ell(L) \leqslant n$. For each idempotent e of M, let $\ell(e)$ denote the maximal length of a P-chain of idempotents ending with e. Then $\ell(e) \leqslant \ell(L)$ by definition. For each $i > 0$, let

$$P_i = \{s \in M \mid \ell(s^\omega) \geqslant i\} \quad \text{and} \quad L_i = \eta^{-1}(P_i)$$

Let $e, f \in E(M)$. Since every idempotent e satisfies $e^\omega = e$, the conditions $e \in P_i$ and $e \leqslant_{\mathcal{J}} f$ imply $f \in P_i$. It follows by Corollary 7.7 that the languages L_i are strongly cyclic. We claim that

$$P = P_1 - P_2 + P_3 - P_4 \ \ldots \ \pm P_m \tag{7.2}$$

First observe that since L is cyclic, an element s of M belongs to P if and only if s^ω belongs to P. Moreover, $s^\omega \in P$ if and only if $\ell(s^\omega)$ is odd. Since $\ell(P) \leqslant n$, one has $\ell(s^\omega) \leqslant n$ for every $s \in M$ and thus $P_{n+1} = \emptyset$. Formula (7.2) follows, since for each $r \geqslant 0$,

$$\{s \in M \mid \ell(s^\omega) = r\} = P_r - P_{r+1}.$$

Moreover, one gets from (7.2) the formula

$$L = L_1 - L_2 + L_3 \ \ldots \ \pm L_n \tag{7.3}$$

which completes the proof of the theorem. □

Theorem 7.10 can be used to give an another proof of Theorem 7.8. To get this result, we must prove that any cyclic language belong to the class $\mathcal{B}_n(\mathcal{S})$ for some integer n. By Theorem 7.10, it suffices to prove that the length of the P-chains of idempotents in a monoid recognising L is bounded. This is a consequence of the following proposition.

Proposition 7.12. *Let L be a regular cyclic language. Let $\varphi : A^* \to M$ be a stamp recognising L and let $P = \varphi(L)$. Then the length of any P-chain of idempotents is bounded by the $\mathcal{J}$-depth of M.*

Proof. Let $(e_0, \ldots, e_{n-1})$ be a P-chain of idempotents in M. Then by definition

$$e_0 \leqslant_{\mathcal{J}} \ \ldots \ \leqslant_{\mathcal{J}} e_{n-1}.$$

Moreover, if $e_{i-1} \ \mathcal{J} \ e_i$, then by [16, Proposition 1.12], the idempotents e_{i-1} and e_i are conjugate. That is, there exist two elements x and y of M such that $xy = e_{k-1}$ and $yx = e_k$. Since L is cyclic, P is also cyclic by Proposition 7.4 and (C_2) implies that $e_{i-1} \in P$ if and only if $e_i \in P$, which contradicts the definition of a P-chain of idempotents. It follows that the sequence $(e_0, \ldots, e_{n-1})$ is a strict $<_{\mathcal{J}}$-chain and hence its length is bounded by the $\mathcal{J}$-depth of M. □

Example 7.3. Let L be the cyclic language $(b+aa)^* + (ab^*a)^* + a^* - b^* + 1$. Its syntactic monoid is the monoid with zero presented by the relations $bb = b$, $a^3 = a$, $baa = a^2b$, $a^2ba = ba$, $bab = 0$. Its transition table and its $\mathcal{J}$-class structure are represented below. The syntactic image of L is $P = \{1, a, a^2, aba, a^2b\}$ and $(aba, b, 1)$ is a maximal P-chain of idempotents.

		1	2	3	4	5	6	7	8
$*$	1	1	2	3	4	5	6	7	8
	a	3	4	5	2	3	8	2	6
$*$	b	7	0	8	4	4	0	7	8
$*$	a^2	5	2	3	4	5	6	4	8
	ab	8	4	4	0	8	8	0	0
	ba	2	0	6	2	2	0	2	6
$*$	a^2b	4	0	8	4	4	0	4	8
$*$	aba	6	2	2	0	6	6	0	0
$*$	bab	0	0	0	0	0	0	0	0



8. Decidability questions on regular languages

Given a lattice of regular languages $\mathcal{L}$, four natural questions arise:

Question 1. Is the membership problem for $\mathcal{L}$ decidable?

Question 2. Is the membership problem for $\mathcal{B}(\mathcal{L})$ decidable?

Question 3. For a given positive integer n, is the membership problem for $\mathcal{B}_n(\mathcal{L})$ decidable?

Question 4. Is the hierarchy $\mathcal{B}_n(\mathcal{L})$ decidable?

More precisely, given a regular language L, Question 1 asks to decide whether $L \in \mathcal{L}$, Question 2 whether $L \in \mathcal{B}(\mathcal{L})$ and Question 3 whether $L \in \mathcal{B}_n(\mathcal{L})$. Question 4 asks whether on can one effectively compute the smallest n such that $L \in \mathcal{B}_n(\mathcal{L})$, if it exists. Note that if Questions 2 and 3 are decidable, then so is Question 4. Indeed, given a language L, one first decides whether L belongs to $\mathcal{B}(\mathcal{L})$ by Question 2. If the answer is positive, this ensures that L belongs to $\mathcal{B}_n(\mathcal{L})$ for some n and Question 3 allows one to find the smallest such n.

If the lattice $\mathcal{L}$ is finite, it is easy to solve the four questions in a positive way. In some cases, a simple application of Corollary 5.5 suffices to solve Question 3 immediately. One just needs to find the appropriate closure operator and to provide algorithms to compute the functions $f(X)$ and $g(X)$ defined by (5.1).

Example 8.1. Let $\mathcal{L}$ be the lattice generated by the languages of the form B^*, where $B \subseteq A$. Then both $\mathcal{L}$ and $\mathcal{B}(\mathcal{L})$ are finite. It is known that a regular language belongs to $\mathcal{L}$ if and only if its ordered syntactic monoid is idempotent and commutative and satisfies the inequation $1 \leqslant x$ for all x. It belongs to $\mathcal{B}(\mathcal{L})$ if and only if its syntactic monoid is idempotent and commutative.

Finally, one can define a closure operator by setting $\overline{L} = B^*$, where B is the set of letters occurring in some word of L. For instance, let $L = \{a, b, c\}^* - \{b, c\}^* + \{a, b\}^* - a^* + 1$. The minimal automaton of L is represented below:



Applying the approximation algorithm of Section 5, one gets $L_0 = \{a, b, c\}^*$, $L_1 = \{b, c\}^*$, $L_2 = b^*$ and $L_3 = \emptyset$ and thus $L = \{a, b, c\}^* - \{b, c\}^* + b^*$ is the best 3-approximation of L.

In the next section, we consider a last series of examples.

9. The difference hierarchy of the polynomial closure of a lattice

A language L of A^* is a *marked product* of the languages $L_0, L_1, \ldots, L_n$ if

$$L = L_0 a_1 L_1 \cdots a_n L_n$$

for some letters $a_1, \ldots, a_n$ of A. Given a set $\mathcal{L}$ of languages, the *polynomial closure* of $\mathcal{L}$ is the set of languages that are finite unions of marked products of languages of $\mathcal{L}$. The *polynomial closure* of $\mathcal{L}$ is denoted Pol $\mathcal{L}$ and the Boolean closure of Pol $\mathcal{L}$ is denoted $\mathscr{B}$Pol $\mathcal{L}$. Finally, let co-Pol $\mathcal{L}$ denote the set of complements of languages in Pol $\mathcal{L}$. In this section, we are interested in the difference hierarchy induced by Pol $\mathcal{L}$. We consider several examples.

9.1. **Shuffle ideals.** If $\mathcal{L} = \{\emptyset, A^*\}$, then Pol $\mathcal{L}$ is exactly the set of shuffle ideals considered in Examples 4.5 and 8.1 and $\mathscr{B}$Pol $\mathcal{L}$ is the class of *piecewise testable languages*. The following easy result was mentioned in [19].

Proposition 9.1. *A language is a shuffle ideal if and only if its syntactic ordered monoid M satisfies the inequation $1 \leqslant x$ for all $x \in M$.*

The syntactic characterisation of piecewise testable languages follows from a much deeper result of Simon [23].

Theorem 9.2. *A language is a piecewise testable if and only if its syntactic monoid is $\mathcal{J}$-trivial.*

Note that the closed sets of the closure operator $X \to X \sqcup A^*$ of Example 4.5. It follows that for the lattice $\mathcal{L}$ of shuffle ideals, the four questions mentioned earlier are decidable. The decidability of Question 3 follows from the approximation algorithm. See Example 5.1.

9.2. **Group languages.** Recall that a *group language* is a language whose syntactic monoid is a group, or, equivalently, is recognized by a finite deterministic automaton in which each letter defines a permutation of the set of states. According to the definition of a polynomial closure, a *polynomial of group languages* is a finite union of languages of the form $L_0a_1L_1\cdots a_kL_k$ where $a_1,\ldots,a_k$ are letters and $L_0,\ldots,L_k$ are group languages.

Let $d_{\mathbf{G}}$ be the metric on A^* defined as follows:

$$r_{\mathbf{G}}(u,v) = \min\{|M| \mid M \text{ is a finite group that separates } u \text{ and } v\}$$
$$d_{\mathbf{G}}(u,v) = 2^{-r_{\mathbf{G}}(u,v)}$$

One can show also that the closure of regular language for $d_{\mathbf{G}}$ is again regular and can be effectively computed [20, 21].

Let $\mathcal{G}$ be the set of group languages on A^* and let Pol $\mathcal{G}$ be the polynomial closure of $\mathcal{G}$. We also let co-Pol $\mathcal{G}$ denote the set of complements of languages of Pol $\mathcal{G}$. The following characterisation of Pol $\mathcal{G}$ was given in [17].

Theorem 9.3. *Let L be a regular language and let M be its ordered syntactic monoid. The following conditions are equivalent:*

(1) *$L \in$ Pol $\mathcal{G}$,*
(2) *L is open in the pro-group topology on A^*,*
(3) *for all $x \in M$, $1 \leqslant x^\omega$.*

Theorem 9.3 shows that Pol $\mathcal{G}$ is decidable. The corresponding result for $\mathscr{B}$Pol $\mathcal{G}$ has a long story, related in detail in [18], where several other characterizations can be found.

Theorem 9.4. *Let L be a regular language and let M be its syntactic monoid. The following conditions are equivalent:*

(1) *$L \in \mathscr{B}$Pol $\mathcal{G}$,*
(2) *the submonoid generated by the idempotents of M is $\mathcal{J}$-trivial,*
(3) *for all idempotents e, f of M, the conditions $efe = e$ implies $ef = e = fe$.*

Theorem 9.4 shows that $\mathscr{B}$Pol $\mathcal{G}$ is decidable. Now, Theorem 9.3 show that a regular language belongs to co-Pol $\mathcal{G}$ if and only if it is closed in the pro-group topology on A^*. It follows that co-Pol $\mathcal{G}$ is closed under arbitrary intersections and the operation associating to a regular language of A^* its closure in the pro-group topology is a closure operator. It follows that the algorithm described at the end of Section 5 can be applied to get our last corollary:

Corollary 9.5. *The difference hierarchy $\mathcal{B}_n($Pol $\mathcal{G})$ is decidable.*

REFERENCES

[1] M.-P. BÉAL, O. CARTON AND C. REUTENAUER, Cyclic languages and strongly cyclic languages, in *STACS '96*, pp. 49–59, *Lect. Notes in Comput. Sci.* vol. 1046, 1996.
[2] J.-Y. CAI, T. GUNDERMANN, J. HARTMANIS, L. A. HEMACHANDRA, V. SEWELSON, K. WAGNER AND G. WECHSUNG, The Boolean hierarchy. I. Structural properties, *SIAM J. Comput.* **17**,6 (1988), 1232–1252.
[3] J.-Y. CAI AND L. HEMACHANDRA, The Boolean hierarchy: hardware over NP, in *Structure in complexity theory (Berkeley, Calif., 1986)*, pp. 105–124, *Lect. Notes in Comput. Sci.* vol. 223, Springer, Berlin, 1986.
[4] O. CARTON, Chain automata, *Theoret. Comput. Sci.* **161** (1996), 191–203.
[5] O. CARTON, A hierarchy of cyclic languages, *R.A.I.R.O.-Informatique Théorique et Applications* **31**,4 (1997), 355–369.

[6] O. CARTON AND D. PERRIN, Chains and superchains in ω-semigroups, in *Semigroups, Automata and Languages*, J. Almeida, G. Gomes and P. Silva (eds.), pp. 17–28, World Scientific, 1994.
[7] O. CARTON AND D. PERRIN, Chains and superchains for ω-rational sets, automata and semigroups, *Int. J. Alg. Comput.* **7**,7 (1997), 673–695.
[8] O. CARTON AND D. PERRIN, The Wadge-Wagner hierarchy of ω-rational sets, in *Automata, Languages and Programming*, P. Degano, R. Gorrieri and A. Marchetti-Spaccamela (eds.), pp. 17–35, *Lect. Notes in Comput. Sci.* vol. 1256, Springer-Verlag, 1997.
[9] O. CARTON AND D. PERRIN, The Wagner hierarchy of ω-rational sets, *Int. J. Alg. Comput.* **9**,5 (1999), 597–620.
[10] C. GLASSER AND H. SCHMITZ, The Boolean structure of dot-depth one, *J. Autom. Lang. Comb.* **6**,4 (2001), 437–452. 2nd Workshop on Descriptional Complexity of Automata, Grammars and Related Structures (London, ON, 2000).
[11] C. GLASSER, H. SCHMITZ AND V. SELIVANOV, Efficient algorithms for membership in Boolean hierarchies of regular languages, *Theoret. Comput. Sci.* **646** (2016), 86–108.
[12] F. HAUSDORFF, *Grundzüge der Mengenlehre. Mit 53 Figuren im Text.*, Leipzig: Veit & Comp., 1914.
[13] F. HAUSDORFF, *Grundzüge der Mengenlehre*, Chelsea Publishing Company, New York, N. Y., 1949.
[14] F. HAUSDORFF, *Set theory*, Chelsea Publishing Company, New York, 1957. Translated by John R. Aumann, et al.
[15] J. KÖBLER, U. SCHÖNING AND K. W. WAGNER, The difference and truth-table hierarchies for NP, *RAIRO Inform. Théor. Appl.* **21**,4 (1987), 419–435.
[16] J.-E. PIN, *Varieties of Formal Languages*, North Oxford Academic, London and Plenum, New York, 1986.
[17] J.-E. PIN, Polynomial closure of group languages and open sets of the Hall topology, in *21th ICALP*, Berlin, 1994, pp. 424–435, *Lect. Notes in Comput. Sci.* n° 820, Springer.
[18] J.-E. PIN, $PG = BG$, a success story, in *NATO Advanced Study Institute Semigroups, Formal Languages and Groups*, J. Fountain (ed.), pp. 33–47, Kluwer academic publishers, 1995.
[19] J.-E. PIN, A variety theorem without complementation, *Russian Mathematics (Iz. VUZ)* **39** (1995), 80–90.
[20] J.-E. PIN AND C. REUTENAUER, A conjecture on the Hall topology for the free group, *Bull. London Math. Soc.* **23** (1991), 356–362.
[21] L. RIBES AND P. A. ZALESSKII, On the profinite topology on a free group, *Bull. London Math. Soc.* **25**,1 (1993), 37–43.
[22] M. P. SCHÜTZENBERGER, Une théorie algébrique du codage, *Séminaire Dubreil. Algèbre et théorie des nombres* **9** (1955-1956), 1–24. http://eudml.org/doc/111094.
[23] I. SIMON, Piecewise testable events, in *Proc. 2nd GI Conf.*, H. Brackage (ed.), pp. 214–222, *Lecture Notes in Comp. Sci.* vol. 33, Springer Verlag, Berlin, Heidelberg, New York, 1975.
[24] W. W. WADGE, Early investigations of the degrees of Borel sets, in *Wadge degrees and projective ordinals. The Cabal Seminar. Volume II*, pp. 166–195, *Lect. Notes Log.* vol. 37, Assoc. Symbol. Logic, La Jolla, CA, 2012.
[25] K. WAGNER, On ω-regular sets, *Inform. and Control* **43**,2 (1979), 123–177.
[26] G. WECHSUNG, On the Boolean closure of NP, in *Fundamentals of computation theory (Cottbus, 1985)*, pp. 485–493, *Lect. Notes in Comput. Sci.* vol. 199, Springer, Berlin, 1985.